

Datasheet

Endopan 3

Special Medium for Endothelial Cells

Product	Description	Catalogue-No.	Size
Endopan 3 ready-to-use	Special formulated complete Medium for endothelial cells	P04-00100	500 ml
Endopan 3 Kit	Special formulated Medium for endothelial cells with 9 Supplements	P04-0010K	500ml

General Information:

Endothelial cells from blood and lymphatic vessels or the internal cavities of the heart play a key role in many physiological and patho-physiological processes. Adhesion among the cells is enabled by desmosomes and tight-junctions. A huge number of soluble factors circulating in the blood or released by neighbouring cells control proliferation or apoptosis of endothelial cells and the invasion of leucocytes to the endothelium, thereby regulating the maintenance, degeneration, or regeneration of blood vessels.

Product Information:

Endopan 3 ready to use is a medium specially developed for the in vitro culture of human endothelial cells containing all components necessary for optimal growth. It is designed for use in an incubator at 37 °C with a 5 % CO₂ atmosphere.

Endopan 3 Kit is provided with FBS and supplements in separate sterile packaging. This will enable the user to prepare a medium for special applications. For example FBS, VEGF, FGF-2 or other components may be omitted from the complete medium for specific experimental settings. Please note that such a formulation will not promote optimal cell growth.

Storage conditions

Storage: Complete and basal medium: 2 – 8 °C in the dark;
Supplements: - 20 °C
Stability: Basal Medium: 12 months
Complete Medium: 3 months
Filling: 500 ml, other sizes on request

Composition

Endopan 3 kit contains the following in single packaging:

- EGF (human recombinant Epidermal Growth Factor)
- FGF-2 (basic Fibroblast Growth Factor)
- VEGF (Vascular Endothelial Growth Factor)
- Vitamin C (Ascorbic acid phosphate)
- R3-IGF-1 (human recombinant Insulin-like Growth Factor)
- FBS (Fetal Bovine Serum)
- G/A (Gentamicin/Amphotericin)
- Hydrocortisone
- Heparin

Suitability

FOR RESEARCH USE ONLY!

Not approved for human or animal diagnostic or therapeutic procedures.

Instructions for use

For a T25 cell culture flask it is recommended to use 5ml of Endopan 3 for smaller or larger culture area, please adjust the volume accordingly.

After thawing cells, change medium after 24 h to remove unattached cells. For maintenance and propagation change medium every two or three days. For cultures close to confluence or for maximum proliferative response it is recommended to use more medium or more frequent medium changes.

The basal and complete medium should not be frozen.

Endopan 3 is a low serum medium (3 % FBS content). It is not suited for stopping trypsin activity. Please use a Trypsin inhibitor to stop the trypsin reaction or use Accutase to detach the cells.

Technical Support

For technical support, questions or remarks please contact your local PAN-Biotech partner or the technical department of PAN-Biotech via email (info@pan-biotech.com) or phone +49-8543-601630.